

7.3 Adaptive responses: T and B cells

Name: _____ Class: _____ Date: _____

Total: 23 marks**KEY WORDS** co-stimulation

Objective

Apply the two-signal rule and compute clonal expansion.

1 Key knowledge and worked examples

■ The gate

Signal 1 alone $\rightarrow$ anergy/deletion; signal 1 + co-stimulation $\rightarrow$ clonal expansion ✓.
CD4 helps, CD8 kills.

■ Worked example: the arithmetic

Doubling ~6-hourly: $2^{12} \approx 4000$ by day 3; $\sim 2^{20} \approx 10^6$ by day 5 ✓. Adjuvant = signal 2.

2 Practice

2.1 What happens to a T cell that gets signal 1 only? [3]

2.2 Name the co-stimulatory pair. [3]

2.3 Which T-cell subset helps B cells class-switch? [3]

3 Exam-style questions

3.1 A vaccine contains purified protein with adjuvant (fictional).

Explain why the adjuvant is necessary (signal 2 via PRR-activated dendritic cells). Predict the response without it. Explain how the two-signal rule prevents autoimmunity to quiet self antigens. [7]

3.2 A boy has recurrent pyogenic infections, high IgM, and absent IgG/IgA (fictional hyper-IgM).

Name the defective molecule (CD40L) and its direction (T $\rightarrow$ B). Explain BOTH infection patterns (no switching $\rightarrow$ no IgG; no macrophage activation $\rightarrow$ opportunists). State the inheritance and why. [7]
