

# Topic 04

## University Medicine

### What this course is, and how to use it

This is the fourth course in University Medicine: **histology and embryology**. MED03 gave the body's map; this course reads its fabric under the microscope and rewinds to how it was woven – the tissue-level structure that explains organ function, and the developmental sequence whose errors predict the anomalies.

**Scope limit, stated once:** a study course, not a medical degree; every case below is fictional and nothing here licenses or guides treatment.

- Units 4.1–4.3 are the tissues: preparation, the four basic types.
- Units 4.4–4.5 are organs read as tissue architecture.
- Units 4.6–4.7 are development: fertilisation to birth, and what failure looks like.

The medicine habit: **tissue architecture is organ function** – every layer, lumen and band below is taught with the physiology it enables and the anomaly its mis-making predicts.

### Microscopy and tissue preparation

**Why it matters.** Histology is not “looking at pictures”: the picture is manufactured, and every step of its manufacture can create or destroy the evidence. Reading a slide means knowing what the preparation did.

**The idea, with a picture.** The sequence: **fixation** (formalin cross-links proteins, preserving structure), **dehydration** (alcohols remove water), **embedding** (paraffin supports the tissue), **sectioning** (microtome, 3–5  $\mu\text{m}$  sections), **staining** (haematoxylin–eosin: H&E, - ).



H = acidic structures blue (nuclei) · E = basic pink (cytoplasm)

*Five steps, each with a failure mode – the artefact you can name is the artefact you can discount*

**Definitions with numbers.** Section thickness 3–5  $\mu\text{m}$  (a cell is 10–30  $\mu\text{m}$  – one

section slices through part of a cell). Resolution of the light microscope  $\sim 200$  nm (half the wavelength of visible light): that limit, not the eye, decides what is visible. Magnification beyond  $\sim 1500\times$  is empty – blur enlarged.

**The derivation (why H&E works).** Haematoxylin is basic and stains acidic structures **blue-purple** (nuclei: DNA and RNA are acids); eosin is acidic and stains basic structures **pink-red** (cytoplasm, collagen). The charge pairing is the whole logic: acid-base electrostatics, not magic – and special stains follow the same rule (PAS stains carbohydrates, so basement membranes and glycogen light up magenta).

**Worked example 1 – the resolution arithmetic.** Two mitochondria 150 nm apart cannot be resolved (below the  $\sim 200$  nm limit) ✓; a 2  $\mu\text{m}$  bacterium and a 10  $\mu\text{m}$  nucleus both can. The limit explains why electron microscopy (wavelength  $\sim 0.005$  nm) exists: it buys  $\sim 1000\times$  the resolving power, at the cost of killing the specimen.

**Worked example 2 – clinical link: the frozen section.** (Fictional vignette) a surgeon waits mid-operation for a margin answer: the frozen-section technique skips dehydration/embedding – the tissue is frozen and cut within minutes, stained quickly, read at lower quality. Speed trades artefact: ice crystals tear the tissue, detail is blurred – but a same-hour answer beats a perfect tomorrow answer.

**Common mistakes.** Treating every gap as a real space (shrinkage during dehydration creates artificial spaces). Reading stain colour as intrinsic colour. Forgetting the section is one plane through a 3-D structure (a tube cut obliquely looks like an oval).

**Summary.** Fix–dehydrate–embed–section–stain, 3–5  $\mu\text{m}$  sections; light microscopy resolves  $\sim 200$  nm; H = acidic blue (nuclei), E = basic pink (cytoplasm); frozen sections trade quality for minutes.

## Epithelium and connective tissue

**Why it matters.** Two of the four tissue types carry the body's whole division of labour: epithelium is the boundary workforce (barrier, transport, secretion), connective tissue is the infrastructure (support, transport highway, immune terrain). Organ function is mostly which epithelium sits on which connective tissue.

**The idea, with a picture. Epithelium** is classified by cell layer (simple one; stratified many) and cell shape (squamous, cuboidal, columnar): simple squamous for exchange (capillary, alveolus), simple cuboidal for transport (tubules), stratified squamous for abrasion (skin, oesophagus). **Connective tissue** = cells + **extracellular matrix** (fibres: collagen, elastic; ground substance), from loose (areolar) to dense regular (tendon) to specialised (bone, blood).

| layer × shape |           |             |            | connective spectrum |               |
|---------------|-----------|-------------|------------|---------------------|---------------|
|               | squamous  | cuboidal    | columnar   | cells + matrix      |               |
| simple        | exchange  | transport   | absorption | loose               | cell wrap     |
|               | capillary | tubule      | gut        | areolar             |               |
| stratified    | abrasion  | wet barrier | protection | dense               | tendon        |
|               | skin      | ureter      | conj.      | regular             |               |
|               |           |             |            | cartilage           | joint         |
|               |           |             |            | bone                | scaffold      |
|               |           |             |            | blood               | plasma matrix |

*Layer × shape classifies every epithelium; fibre density and arrangement classifies every connective tissue*

**The derivation (structure follows function).** Exchange needs minimal diffusion distance: one flat layer (simple squamous, ~0.5 µm) ✓. Abrasion defence needs replaceable depth: many layers (stratified squamous) ✓. Transport and secretion need machinery: tall cells packed with organelles (cuboidal/columnar). The classification grid is not taxonomy for its own sake – each cell of the grid predicts the underlying physiology.

**Worked example 1 – reading a slide.** A section shows a free surface of tall cells with nuclei in a basal row, on a thin basement membrane, over loose matrix with small vessels: simple columnar epithelium on areolar connective tissue – a transport/absorption surface (intestine pattern). Every clue (apical surface, basal membrane, matrix) is a coordinate in the grid.

**Worked example 2 – clinical link: the barrier that fails.** (Fictional biopsy) recurrent infections, steatorrhoea: the duodenal biopsy shows flattened villi – the absorptive simple columnar epithelium’s surface area has collapsed. Malabsorption is not a “functional” mystery: the exchange architecture itself is gone, and the microscope reads the mechanism.


**Common mistakes.** Confusing the basement membrane (epithelium’s anchor) with the whole matrix. Calling blood “not connective tissue” (it is: cells + matrix = plasma). Forgetting pseudostratified looks layered but IS one layer (all cells touch the basement membrane – respiratory epithelium).

**Summary.** Layer × shape classifies epithelium; each grid cell predicts function (exchange = simple squamous; abrasion = stratified); connective tissue = cells + collagen/elastic/ground substance, from areolar to tendon to bone and blood.

## Muscle and nervous tissue

**Why it matters.** Movement and control are tissues with visible machinery: the sarcomere's banding IS the contraction mechanism, and the neuron's shape IS its connectivity. Histology here is physiology you can see.

**The idea, with a picture. Muscle** : skeletal (striated, voluntary, multi-nucleate at the periphery), cardiac (striated, involuntary, branched with intercalated discs), smooth (unstriated, involuntary, spindle cells). **Nervous tissue** : neurons (cell body, dendrites, axon) plus **glia** (astrocytes support, oligodendrocytes/Schwann cells myelinate, microglia scavenge).



*A-band stays constant (myosin); the I-band and H-zone shrink as the filaments slide – the bands are the mechanism*

**The sarcomere** derivation. A **sarcomere** runs Z-line to Z-line. The **A-band** (anisotropic, dark) is myosin's full length – it does NOT change in contraction. The **I-band** (isotropic, light) is actin-only; the **H-zone** is myosin-only within the A-band. As actin slides toward the centre, the I-band and H-zone NARROW and the Z-lines approach – filament lengths never change ✓. The banding you see is the sliding-filament model printed on the tissue.

**Worked example 1 – identifying muscle in a section.** Peripheral nuclei + long parallel fibres + striations = skeletal ✓. Central nuclei, branching, intercalated discs = cardiac ✓. No striations, spindle cells, central single nuclei = smooth ✓. Three tissues, three fingerprints, no special stains needed.

**Worked example 2 – clinical link: myelin read under the scope.** (Fictional nerve biopsy) a child with progressive weakness shows “onion bulbs” – concentric Schwann-cell processes around thinly myelinated axons: repeated de/remyelination made visible. The histology names the disease class (demyelinating neuropathy) because glia are as much “nervous tissue” as neurons.

**Common mistakes.** Expecting the A-band to shorten (it is myosin's constant length). Confusing intercalated discs with striations (discs are junctions between cells, dark lines across the fibre). Treating glia as packing material (they myelinate, buffer,


and scar).

**Summary.** Three muscle fingerprints (peripheral-nuclei striated / branched with discs / unstriated spindle); sarcomere: A constant, I and H narrow, filaments never shorten; glia are functional nervous tissue – myelin pathology is glial pathology.

## Organ histology: circulatory, respiratory and digestive

**Why it matters.** An organ is a tissue arrangement solving a physiological problem – and the three great tube systems (vessels, airways, gut) wear their haemodynamics on their walls. Read the layers and you read the pressures.

**The idea, with a picture.** Vessels: **tunica intima** (endothelium), **media** (smooth muscle – thick in arteries, thin in veins), **adventitia** (collagen). Airway: the wall thins from bronchus to **alveolus** where the wall becomes simple squamous over capillaries – the blood–air barrier  $\sim 0.5\ \mu\text{m}$  ✓. Gut: **mucosa** (epithelium + lamina propria + muscularis mucosae), **submucosa**, **muscularis externa** (inner circular, outer longitudinal), **serosa** or adventitia.



artery = mostly media (pulse) · vein = thin walls · gut = 4 coats

*Three tubes, one grammar: lining, muscle coat, outer sheath – each layer named for its job*

**The derivation (layers track pressure and traffic).** Arteries carry high-pressure pulse: a thick elastic/muscular media smooths it ✓. Veins carry low-pressure volume: thin walls, valves, big lumen ✓. The alveolus trades every defence for distance: no muscle, no glands, one flat cell plus its basement membrane and the capillary wall – exchange optimised to the exclusion of everything else ✓. The gut keeps a muscularis externa because its job is propulsion, plus a mucosa that is the body's most renewable frontline.

**Worked example 1 – artery vs vein at a glance.** The vessel with the thick media nearly filling the wall and a round lumen is the artery; the floppy wide-lumen vessel with a thin wall is the vein ✓. The histological look IS the pressure record – an artery is "mostly media", a vein is "mostly lumen".

**Worked example 2 – clinical link: the barrier that decides gas exchange.** (Fictional case) breathless with clear airways: the alveolar wall has thickened (fibrosis pattern) – the 0.5  $\mu\text{m}$  barrier has become several  $\mu\text{m}$ , and diffusion length, not airflow, now limits oxygen. The histology explains why the inhaler cannot help: the problem is in the wall, not the tube.

**Common mistakes.** Calling every inner layer “mucosa” (vessels have intima; mucosa belongs to wet epithelial tubes). Forgetting the gut’s two muscle coats run in opposite directions (circular squeezers, longitudinal chasers). Expecting alveoli to look like bronchi (they are the opposite: everything removed).

**Summary.** Vessel = intima/media/adventitia, media  $\propto$  pressure; gut = mucosa/submucosa/muscularis externa/serosa; alveolus = simple squamous


- basement membrane + capillary,  $\sim 0.5 \mu\text{m}$  total; the layer list is the pressure/traffic profile.

## Organ histology: urinary, endocrine, reproductive and skin

**Why it matters.** The kidney’s tubule segments, the endocrine’s vascular nets, the skin’s stratified armour – each is a recognisable histological machine, and together they cover the urine, the hormones and the barrier: three systems whose diseases are read in biopsies every day.

**The idea, with a picture.** The **nephron** : glomerulus (fenestrated capillary tuft in Bowman’s capsule)  $\rightarrow$  proximal tubule (cuboidal with brush border)  $\rightarrow$  loop of Henle (flat squamous thin limbs, cuboidal thick limb)  $\rightarrow$  distal tubule (cuboidal, no brush)  $\rightarrow$  collecting duct (cuboidal to columnar, with pale principal cells). Endocrine organs = cords or follicles wrapped in **fenestrated capillaries** (secreting into blood needs the shortest path). Skin = **stratified squamous keratinised** on dermis, with the epidermis turning over every  $\sim 4$  weeks.

filtrate flows down



*Each segment wears its job: brush border for reabsorption, flat for exchange, columnar for ducts*

**The derivation (epithelium predicts tubule function).** The proximal tubule reabsorbs ~65 % of the filtrate: it needs surface area – hence a **brush border** (microvilli) that stains brightly ✓. The loop of Henle’s thin limbs let water and ions move passively: flat squamous, minimal machinery ✓; its thick ascending limb pumps salt out and is cuboidal, packed with mitochondria ✓. The collecting duct conveys and fine-tunes: cuboidal to columnar, with water channels under hormonal control ✓. You can deduce a segment’s physiology from one high-power field.

**Worked example 1 – biopsy orientation.** A field shows a tuft of capillaries inside a capsule, beside a tubule with a fuzzy violet brush border: glomerulus plus proximal tubule ✓ – renal cortex. The signature pairing names the region without a map.

**Worked example 2 – clinical link: the four-week frontline.** (Fictional burn vignette) the barrier is lost where the keratinising epithelium died: fluid loss and infection follow immediately, and healing depends on how much **basement membrane** survived, with the hair follicles and glands below it (they are where new epidermis grows from). One layer’s survival decides the prognosis – histology as triage.


**Common mistakes.** Expecting every kidney tubule to look the same (the segments are the point). Forgetting endocrine cells sit directly ON capillaries (no ducts – that is what “endocrine” means). Calling the dermis “just connective tissue” (it is the vascular, sensory, structural half of skin).

**Summary.** Nephron segments read as epithelial signatures (brush = proximal; flat = thin loop; cuboidal to columnar = collecting); endocrine = secretory cells on fenestrated capillaries; skin = keratinised stratified squamous over vascular dermis, ~4-week turnover; basement membrane survival decides regeneration.

## Early development: fertilisation to gastrulation

**Why it matters.** The first three weeks build the body’s axes and layers from one cell – with a speed and precision that make them both the most vulnerable and the most instructive period of life. Every later organ trace begins here.

**The idea, with a picture.** Timeline: **fertilisation** (day 0) → **cleavage** → **morula** (~day 3) → **blastocyst** (~day 5, implants ~day 6–7) → **gastrulation** (week 3: forms the three germ layers) → **neurulation** (week 3–4: the neural tube closes).



*Day 0 to week 3: one cell becomes three layers – the whole body is a mapping of these three*

**The derivation (why gastrulation is the pivot).** The blastocyst is a ball of cells; the body is three layered **germ layers** with axes – **ectoderm** (nervous system, epidermis), **mesoderm** (muscle, bone, blood, kidney, heart), **endoderm** (gut, lungs' lining, liver, pancreas) ✓. Gastrulation is the reorganisation that CREATES the layers and the axes; everything after is specialisation. The germ-layer map is why a single week-3 field defect predicts a multi-organ syndrome: tissues are neighbours before they are organs.

**Worked example 1 – the timeline arithmetic.** Fertilisation to implantation  $\approx$  6–7 days ✓; to gastrulation  $\approx$  15 days; to neural-tube closure completion  $\approx$  day 28. Why folate matters periconceptionally: the neural tube closes in week 4 after conception – about two weeks after the first missed period, often before the pregnancy is known or the first antenatal visit – so the protective supplement must start before conception ✓.

**Worked example 2 – clinical link: the tube that fails to close.** (Fictional case) elevated maternal alpha-fetoprotein, ultrasound shows an open spine: neural-tube closure failed at day 26–28. The defect level maps the deficit (lumbar → legs and bladder; cervical → quadriplegia) – development wrote the prognosis in anatomy.

**Common mistakes.** Mixing up the blastocyst's two parts (the inner cell mass becomes the embryo; the trophoblast becomes the placenta). Treating the germ layers as a list to memorise (they are a spatial map: tube-in-tube-in-body). Counting pregnancy weeks from conception (clinical weeks count from the last period:  $\sim$ 2 weeks ahead).


**Summary.** Day 0 fertilisation → day 5 blastocyst → week 3 gastrulation makes ectoderm/mesoderm/endoderm → week 4 neural tube; the germ-layer map predicts organ systems and syndrome patterns; closure is complete about two weeks after the

first missed period, so folate must start before conception.

## Organogenesis and congenital anomalies

**Why it matters.** Weeks 4–8 turn three layers into recognisable organs – the most error-prone window in human life. Congenital anomaly reasoning is developmental logic read backwards: from a defect, infer the week and the process that failed.

**The idea, with a picture.** Heart septation : the heart tube folds (week 4), then septa close the atrial and ventricular holes – the atrial septum's **foramen ovale** must stay open in the fetus (high pulmonary resistance) and close functionally after birth ✓. Meanwhile: the palate's shelves fuse (weeks 7–10), the diaphragm closes (week 7), the limb buds pattern out (weeks 4–8).



*The most error-prone construction project in the body: each septum has its week, and its failure has its name*

**The derivation (the timing predicts the defect).** Organs form in overlapping windows; a teratogen or gene error hits whatever is mid-construction AT THAT WEEK. Weeks 3–4 (neural tube, heart tube): the classic malformation window; weeks 4–8 (septation, palate, limbs): the structural window; after week 8, growth and function mature rather than form ✓. Anomaly reasoning = matching defect to window: defects in organs that were forming in the same weeks (heart septa, palate, limbs) can share one cause, so a cluster is a clue, not a coincidence.

**Worked example 1 – the septal arithmetic.** The fetal circulation bypasses the lungs: right atrium → foramen ovale → left atrium, and pulmonary artery → ductus arteriosus → aorta ✓. At birth the first breath drops pulmonary resistance, left atrial pressure rises, and the foramen slams shut (functionally immediately, anatomically in months). The ductus arteriosus normally constricts over the first 24–72 hours in term infants; anatomical closure follows later. Patent foramen ovale (persistent in ~25 % of adults) usually matters only when something else forces a right-to-left shunt ✓.

**Worked example 2 – clinical link: one cause, several systems.** (Fictional child) cleft palate and a heart murmur from a ventricular septal defect. The palate and the ventricular septum were both being built in weeks 4–10, and some causes hit both at once: a teratogen in that window, or a genetic change in a signal both use (the 22q11 deletion is the classic one). So one anomaly is a reason to look for its neighbours: the doctor examining a child with a cleft always listens to the heart.

**Common mistakes.** Treating “congenital” as “genetic” (most anomalies are multifactorial; many are pure timing). Dating anomalies by birth findings (the error happened weeks earlier). Forgetting the placenta is fetal tissue (its vessels are part of fetal cardiovascular design).

**Summary.** Weeks 4–8 organogenesis: heart folds then septates, palate fuses, limbs pattern; defects date to their construction windows; the fetal foramen ovale and ductus must be open in utero and close functionally after birth; organs built in the same weeks can share one cause.