

Exercise walk-through

1.6

Cell signalling


University Medicine

Signalling: amplify and reset


- 1 **Ledger** Resting cAMP $\approx 12\,000$ molecules per hepatocyte; ten cyclases at $1000/s$ replace the pool in a second - OFF matters as much as ON.
- 2 **Switch** cAMP switches $10\text{ nM} \rightarrow 1\text{ }\mu\text{M}$ (a hundredfold) in seconds - a maximal hormone pulse.
- 3 **Welded** Cholera toxin disables the $G\alpha_s$ GTPase: cyclase runs constitutively, Cl^- and water pour out - the OFF enzyme IS the story.

Two cascades

GPCR — second messenger



RTK — kinase cascade



GPCR-cAMP and RTK-RAS side by side: $\sim 10^4$ amplification, every stage resettable.

Exercises 2.1-2.3

2.1 Name the OFF timer of the $G\alpha$ subunit. [3]

2.2 Compute the cAMP fold-change from 10 nM to 1 μ M. [3]

2.3 Why do GPCR ligands not enter the cell? [3]

Exercises 2.1-2.3

2.1 Name the OFF timer of the $G\alpha$ subunit. [3]

2.1

- Its intrinsic GTPase: $GTP \rightarrow GDP$, seconds.

2.2 Compute the cAMP fold-change from 10 nM to 1 μ M. [3]

2.2

- $10^{-6}/10^{-8} = 100\times$.

2.3 Why do GPCR ligands not enter the cell? [3]

2.3

- The message is relayed by the transmembrane receptor and second messengers; hydrophilic ligands cannot cross the bilayer (only lipophilic steroids enter, by a different mechanism).

Exercise 3.1

3.1 A maximal hormone pulse.

Compute the cAMP molecules at 1 μM in a 2×10^{-12} L cell (compare the 12 000 resting). Explain the amplification chain that produces them. State two OFF mechanisms and their timescale. [8]

Exercise 3.1

3.1 A maximal hormone pulse.

Compute the cAMP molecules at 1 μM in a 2×10^{-12} L cell (compare the 12 000 resting). Explain the amplification chain that produces them. State two OFF mechanisms and their timescale. [8]

3.1

- $10^{-6} \times 2 \times 10^{-12} \times N_A \approx 1.2 \times 10^6$ molecules – 100× the 12 000 resting ✓.
- Hormone $\rightarrow$ receptor $\rightarrow$ ~ 20 Gs $\rightarrow$ cyclases $\rightarrow$ cAMP: stage gains multiply ($\sim 10^4$).
- OFF: $G\alpha$ GTPase (seconds); PDE destroying cAMP (seconds).

Exercise 3.2

3.2 Cholera toxin (fictional vignette for the mechanism).

State the molecular modification and its direct effect on $G\alpha_s$. Trace the cascade to luminal Cl^- and water loss. Explain why therapy is fluid replacement rather than an inhibitor.

[7]

Exercise 3.2

3.2 Cholera toxin (fictional vignette for the mechanism).

State the molecular modification and its direct effect on $G\alpha_s$. Trace the cascade to luminal Cl^- and water loss. Explain why therapy is fluid replacement rather than an inhibitor. [7]

3.2

- ADP-ribosylation of $G\alpha_s$; GTPase disabled $\rightarrow$ Gs permanently on.
- Constitutive cyclase $\rightarrow$ maximal cAMP $\rightarrow$ PKA $\rightarrow$ Cl^- channel open $\rightarrow$ Cl^- and water follow into the lumen.
- No inhibitor can unweld an already-modified protein pool; support while proteins turn over.