

Exercise walk-through

7.3


Adaptive responses: T and B cells

University Medicine

Adaptive responses: the gate and the arithmetic

- 1 Division** CD4 helps (macrophages, B switching via CD40L); CD8 kills (granules, Fas); B → plasma (thousands of Ig/second) + memory.
- 2 Gate** Signal 1 alone → anergy/deletion; signal 1 + **co-stimulation** (B7–CD28) → clonal expansion - the tolerance/activation decision.
- 3 Arithmetic** One naive cell doubling ~6-hourly: $2^{12} \approx 4000$ by day 3, $\sim 10^6$ by day 5; CD8 >1000-fold of the specific pool.

Adaptive responses - figure



Signal 1 + **co-stimulation**: tolerise or activate.

Exercises 2.1-2.3

2.1 What happens to a T cell that gets signal 1 only? [3]

2.2 Name the co-stimulatory pair. [3]

2.3 Which T-cell subset helps B cells class-switch? [3]

Exercises 2.1-2.3

2.1 What happens to a T cell that gets signal 1 only? [3]

2.1

- Anergy or deletion – tolerance.

2.2 Name the co-stimulatory pair. [3]

2.2

- B7 (dendritic) – CD28 (T cell) ✓.

2.3 Which T-cell subset helps B cells class-switch? [3]

2.3

- CD4 (TH) via CD40L–CD40 contact ✓.

Exercise 3.1

3.1 A vaccine contains purified protein with adjuvant (fictional).

Explain why the adjuvant is necessary (signal 2 via PRR-activated dendritic cells). Predict the response without it. Explain how the two-signal rule prevents autoimmunity to quiet self antigens.

[7]

Exercise 3.1

3.1 A vaccine contains purified protein with adjuvant (fictional).

Explain why the adjuvant is necessary (signal 2 via PRR-activated dendritic cells). Predict the response without it. Explain how the two-signal rule prevents autoimmunity to quiet self antigens. [7]

3.1

- Protein alone gives signal 1 without PRR-driven B7 – tolerance or a weak response.
- Without adjuvant: little/no antibody, possible anergy ✓.
- Quiet self-presentation never carries B7, so self-reactive clones are tolerised – autoimmunity needs danger to license them.

Exercise 3.2

3.2 A boy has recurrent pyogenic infections, high IgM, and absent IgG/IgA (fictional hyper-IgM).

Name the defective molecule (CD40L) and its direction (T→B). Explain BOTH infection patterns (no switching → no IgG; no macrophage activation → opportunists). State the inheritance and why. [7]

Exercise 3.2

3.2 A boy has recurrent pyogenic infections, high IgM, and absent IgG/IgA (fictional hyper-IgM).

Name the defective molecule (CD40L) and its direction (T→B). Explain BOTH infection patterns (no switching → no IgG; no macrophage activation → opportunists). State the inheritance and why. [7]

3.2

- CD40L (CD154) on T cells, missing – X-linked.
- No CD40 signal: B cells cannot switch (IgM piles) → pyogenic (encapsulated) infections; macrophages are not activated → opportunists (Pneumocystis, Cryptosporidium).
- X-linked recessive: mother carrier, half of sons affected ✓.