

## Solutions

---

### 2.1

- Its intrinsic GTPase:  $\text{GTP} \rightarrow \text{GDP}$ , seconds.

### 2.2

- $10^{-6}/10^{-8} = 100\times$ .

### 2.3

- The message is relayed by the transmembrane receptor and second messengers; hydrophilic ligands cannot cross the bilayer (only lipophilic steroids enter, by a different mechanism).

### 3.1

- $10^{-6} \times 2 \times 10^{-12} \times N_A \approx 1.2 \times 10^6$  molecules –  $100\times$  the 12 000 resting ✓.
- Hormone  $\rightarrow$  receptor  $\rightarrow \sim 20$  Gs  $\rightarrow$  cyclases  $\rightarrow$  cAMP: stage gains multiply ( $\sim 10^4$ ).
- OFF:  $\text{G}\alpha$  GTPase (seconds); PDE destroying cAMP (seconds).

### 3.2

- ADP-ribosylation of  $\text{G}\alpha$ s; GTPase disabled  $\rightarrow$  Gs permanently on.
- Constitutive cyclase  $\rightarrow$  maximal cAMP  $\rightarrow$  PKA  $\rightarrow$   $\text{Cl}^-$  channel open  $\rightarrow$   $\text{Cl}^-$  and water follow into the lumen.
- No inhibitor can unweld an already-modified protein pool; support while proteins turn over.