

Solutions

2.1

- Muscle lacks glucose-6-phosphatase, so G-6-P cannot become free glucose for export.

2.2

- Glucagon; liver only (no muscle receptors).

2.3

- $4 \text{ ATP} + 2 \text{ GTP} = 6 \text{ high-energy phosphates (plus 2 NADH)}$.

3.1

- Hours 0–4 absorbed fuels; 4–12 hepatic glycogenolysis; 12–24 gluconeogenesis dominant; day 3+ ketones cover $\sim 2/3$ brain needs.
- $100 \text{ g} \div \sim 5 \text{ g/h} \approx \text{up to 24 h}$; strictly, glycogen alone ≈ 1 day of brain use (400 vs 480 kcal).
- Substrates: alanine (muscle), glycerol (fat), lactate (Cori).
- Ketones substitute for glucose $\rightarrow$ less gluconeogenesis $\rightarrow$ protein sparing.

3.2

- Muscle glycogen phosphorylase.
- No glycogen $\rightarrow$ no extra G-6-P flux $\rightarrow$ no pyruvate surge $\rightarrow$ no lactate.
- PFK-1 deficiency: glycogen BREAKS down but flux dies at the gate – also low lactate, but with phosphate intermediates accumulating.
- Advice: aerobic, low-intensity exercise (fat/fatty acids still usable); avoid sprints.