

Solutions

2.1

- Insulin; drives hepatic glycogen synthesis (and lipogenesis).

2.2

- Ketones supply $\sim 2/3$ of brain energy (glucose need falls to $\sim 30\text{--}40$ g/day).

2.3

- Non-enzymatic glycation of haemoglobin, integrated over the ~ 120 -day red-cell lifetime.

3.1

- Hour 6: liver on glycogenolysis, fat basal release, muscle glucose/ fatty acids. Hour 18: glycogen near-empty – gluconeogenesis from amino acids/glycerol/lactate; fat lipolysis high. Day 4: liver exports ketones; muscle largely fatty acids; brain $\sim 2/3$ ketones.
- Switch justified: $100\text{ g bank} \div \sim 120\text{ g/day brain demand} \approx \text{one day}$.
- Protein sparing: each gram of brain glucose replaced by ketones spares ~ 2 g muscle protein (amino-acid carbon for glucose).

3.2

- No insulin $\Rightarrow$ glucagon programme runs: lipolysis (free fatty acids $\rightarrow$ ketones $\rightarrow$ acidosis), proteolysis (weight loss), glycogenolysis + gluconeogenesis despite high glucose; glucose above renal threshold $\rightarrow$ osmotic diuresis (polyuria, thirst); acidosis drives Kussmaul (deep) breathing.
- Type 2: insulin present but resisted – same ledger, milder arrows, years not hours.