

1. Haemoglobin has a quaternary structure but myoglobin does not. Why?

- ☐ Haemoglobin has no α -helices
- ☐ Haemoglobin has four chains; myoglobin has one
- ☐ Myoglobin has no haem group
- ☐ Myoglobin is not a protein

2. An enzyme makes a reaction go further towards products at equilibrium.

- ☐ True
- ☐ False

3. A reaction has $\Delta G^\circ = +14.2$ kJ per mol. It is coupled to ATP hydrolysis (-30.5 kJ per mol). What is the overall ΔG° , in kJ per mol?

_____ kJ per mol

4. One glucose gives two acetyl-CoA. How many NADH do the two TCA turns make?

_____ NADH

5. Which glycogen store keeps blood glucose up between meals?

- ☐ Both equally
- ☐ Liver glycogen
- ☐ Brain glycogen
- ☐ Muscle glycogen

6. How many β -oxidation cycles does stearate (C18) need?

_____ cycles

7. Alanine transaminates with α -ketoglutarate. What are the products?

- ☐ Aspartate and oxaloacetate
- ☐ Urea and water
- ☐ Glucose and ammonia
- ☐ Pyruvate and glutamate

8. How does glucagon turn on glycogen breakdown in the liver?

- ☐ It breaks glycogen down directly
- ☐ It removes phosphates from the enzymes
- ☐ It enters the nucleus and switches on genes within seconds

9. Myoglobin binds O_2 cooperatively, giving a sigmoid curve.

- ☐ True
- ☐ False

10. What is K_m ?

- ☐ The substrate concentration at half of V_{max}
- ☐ The rate when $[S] = 1$ mM
- ☐ The maximum rate of the enzyme
- ☐ The amount of enzyme in the cell

1. **Haemoglobin has four chains; myoglobin has one**

Quaternary structure is the assembly of several chains. One chain (myoglobin) cannot have it; Hb's $\alpha_2\beta_2$ tetramer does.

2. **False**

An enzyme speeds both directions equally; it reaches the same equilibrium sooner but does not move it.

3. **-16.3 kJ per mol**

$+14.2 + (-30.5) = -16.3$ kJ per mol: negative, so the coupled pair runs.

4. **6 NADH**

3 NADH per turn $\times$ 2 turns = 6, plus 2 FADH₂ and 2 GTP.

5. **Liver glycogen**

The liver's ~100 g is exported as glucose; muscle's larger ~400 g is used only by the muscle itself.

6. **8 cycles**

$18/2 - 1 = 8$ cycles, giving 9 acetyl-CoA: the last cycle splits a 4-carbon piece into two acetyl-CoA.

7. **Pyruvate and glutamate**

The amino group moves to α -ketoglutarate, making glutamate; alanine's carbon skeleton is left as pyruvate.

8. **It raises cAMP, and PKA phosphorylates the enzymes**

Glucagon $\rightarrow$ cAMP $\rightarrow$ PKA $\rightarrow$ phosphorylase on, glycogen synthase off. Insulin does the opposite, by removing those phosphates.

9. **False**

Myoglobin has one site, so its curve is a hyperbola. Cooperativity needs several sites that influence each other, as in Hb.

10. **The substrate concentration at half of V_{max}**

Put $[S] = K_m$ into the equation: $v = V_{max}K_m/2K_m = V_{max}/2$.